

Tuning of Monte Carlo Event Generator PYTHIA 6 to Min Bias Data (Preliminary Results)

Nameeqa Firdous, Gerald Rudolph

Institute of Astro and Particle Physics Innsbruck, Austria

Sun July 18, 2010



Motivation

- Min Bias is a cocktail of hard (pQCD) and soft processes



Motivation

- Min Bias is a cocktail of hard (pQCD) and soft processes
- Soft interactions are the largest but most difficult part of the inelastic Xs at Colliders



Motivation

- Min Bias is a cocktail of hard (pQCD) and soft processes
- Soft interactions are the largest but most difficult part of the inelastic Xs at Colliders
- Understand the whole event (not just the pQCD part)



Motivation

- Min Bias is a cocktail of hard (pQCD) and soft processes
- Soft interactions are the largest but most difficult part of the inelastic Xs at Colliders
- Understand the whole event (not just the pQCD part)
- No MC generator can reproduce all MB observables at the same time



Motivation

- Min Bias is a cocktail of hard (pQCD) and soft processes
- Soft interactions are the largest but most difficult part of the inelastic Xs at Colliders
- Understand the whole event (not just the pQCD part)
- No MC generator can reproduce all MB observables at the same time
- To describe data the best possible way **MC models need to be tuned**



PYTHIA Model

- Multiparton Interaction Model
- Pt ordered parton shower



PYTHIA Model

- Multiparton Interaction Model
- Pt ordered parton shower
- “In the full event generation process, probably no other area is as poorly understood as this one!”
(PYTHIA manual)



MPI Model

- Several parton pairs can undergo (semi)hard interaction in an event



MPI Model

- Several parton pairs can undergo (semi)hard interaction in an event
- The hard scatter cross section is extended to the low p_t (diverges for $p_t \rightarrow 0$)



MPI Model

- Several parton pairs can undergo (semi)hard interaction in an event
- The hard scatter cross section is extended to the low p_t (diverges for $p_t \rightarrow 0$)
- A regularization parameter is introduced



MPI Model

- Several parton pairs can undergo (semi)hard interaction in an event
- The hard scatter cross section is extended to the low p_t (diverges for $p_t \rightarrow 0$)
- A regularization parameter is introduced
- All interactions are simulated in decreasing order of p_t



Strategy

- Fragmentation Parameters were tuned to e^+e^- data(ALEPH)



Strategy

- Fragmentation Parameters were tuned to e^+e^- data(ALEPH)
- Multiparton Interaction model parameters are tuned to pp/pp^- data



Fit Procedure

- Linear parameterisation is used, most efficient with the possibility of iteration, allows many (10-20) parameters to vary simultaneously



Fit Procedure

- Linear parameterisation is used, most efficient with the possibility of iteration, allows many (10-20) parameters to vary simultaneously
- Central value and tuning interval for each parameter are chosen, and run the generator with these settings



Fit Procedure

- Linear parameterisation is used, most efficient with the possibility of iteration, allows many (10-20) parameters to vary simultaneously
- Central value and tuning interval for each parameter are chosen, and run the generator with these settings
- This prediction is fitted to data (minimal χ^2)



List of observables

Analysis	Observable	Analysis	Observable
ATLAS 0.9TeV, $n_{ch} \geq 1$ $p_t > 0.5\text{GeV}$ $ \eta < 2.5$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{d\eta}$	CDF-II 1.96TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{dn_{ch}}$
ATLAS 0.9TeV, $n_{ch} \geq 1$ $p_t > 0.5\text{GeV}$ $ \eta < 2.5$	$\frac{1}{N_{ev}} \cdot \frac{1}{2\pi p_t} \cdot \frac{d^2 N_{ch}}{d\eta dp_t}$	CDF-II 1.96TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\frac{1}{N_{ev}} \cdot \frac{1}{2\pi p_t} \cdot \frac{d^2 N_{ch}}{dy dp_t}$
ATLAS 0.9TeV, $n_{ch} \geq 1$ $p_t > 0.5\text{GeV}$ $ \eta < 2.5$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{dn_{ch}}$	CDF-II 1.96TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\frac{1}{N_{ev}} \cdot \frac{1}{2\pi} \cdot \frac{d^2 N_{ch}}{d\eta dE_t}$
ATLAS 0.9TeV, $n_{ch} \geq 1$ $p_t > 0.5\text{GeV}$ $ \eta < 2.5$	$\langle p_t \rangle \text{ vs. } n_{ch}$	CDF-II 1.96TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\langle p_t \rangle \text{ vs. } n_{ch}$
CMS 7TeV, no p_t cut $ \eta < 2.4$	$\frac{1}{N_{ev}} \cdot \frac{1}{2\pi p_t} \cdot \frac{d^2 N_{ch}}{d\eta dp_t}$	CDF-I 0.63TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{dn_{ch}}$
ALICE 7 TeV, no p_t cut, $n_{ch} \geq 1$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{dn_{ch}}$	CDF-I 1.8TeV, $n_{ch} \geq 1$ $p_t > 0.4\text{GeV}$ $ \eta \leq 1.0$	$\frac{1}{N_{ev}} \cdot \frac{dN_{ev}}{dn_{ch}}$

ALICE 0.9 TeV, no p_t cut, $n_{ch} \geq 1$	$\left. \frac{dn_{ch}}{d\eta} \right _{\eta=0}$
ALICE 7 TeV, no p_t cut, $n_{ch} \geq 1$	$\left. \frac{dn_{ch}}{d\eta} \right _{\eta=0}$
ATLAS 0.9 TeV, $p_t > 0.5\text{GeV}$, $n_{ch} \geq 6$	$\left. \frac{dn_{ch}}{d\eta} \right _{\eta=0}$
ATLAS 7TeV, $p_t > 0.5\text{GeV}$, $n_{ch} \geq 6$	$\left. \frac{dn_{ch}}{d\eta} \right _{\eta=0}$
ATLAS 7TeV, $p_t > 0.5\text{GeV}$, $n_{ch} \geq 1$	$\left. \frac{dn_{ch}}{d\eta} \right _{\eta=0}$



List of tuned parameters

Parameters	Pythia Default	Tune CTEQ5			Tune MRSTLO*			Description
		CDF	LHC	Both	CDF	LHC	Both	
PARP(82)	2.00	1.78	1.69	1.76	2.01	1.92	1.99	P10 at reference Ecm
PARP(83)	1.8	2.09	2.54	2.24	2.18	2.37	2.27	matter distribution
PARP(78)	0.03	0.49	0.52	0.51	0.39	0.47	0.41	colour reconnection in FS
PARP(90)	0.16	0.22	0.23	0.24	0.21	0.22	0.22	p10 evolution with Ecm
PARP(77)	0	1.64	1.64	1.64	1.64	1.64	1.64	Suppression of CR for fast moving strings



	Tune CTEQ5				Tune MRSTLO*			AMBT1		
	CDF	LHC	Both		CDF	LHC	Both	CDF	LHC	Both
Chi ² /dof	7.16	2.00	5.23	14.10	3.42	11.38	7.01	1.58	6.01	

Central Charged Densities



Analysis				Data	MC Overall fit	
ALICE	0.9 TeV,	no p_t cut,	$n_{ch} \geq 1$	3.81 ± 0.07	3.56	3.60
ALICE	7 TeV,	no p_t cut,	$n_{ch} \geq 1$	6.01 ± 0.15	5.18	4.81
ATLAS	0.9 TeV,	$p_t > 0.5 \text{ GeV}$,	$n_{ch} \geq 6$	2.38 ± 0.08	2.43	2.45
ATLAS	7 TeV,	$p_t > 0.5 \text{ GeV}$,	$n_{ch} \geq 6$	3.64 ± 0.12	3.69	3.40
ATLAS	7 TeV,	$p_t > 0.5 \text{ GeV}$,	$n_{ch} \geq 1$	2.42 ± 0.08	2.26	2.34





N_ch, MB, 0.63 TeV, CDF-1





N_ch, MB, 1.8 TeV, CDF-1





N_ch, MB, 1.96 TeV, CDF-2





N_ch, MB, 0.9 TeV, ATLAS








pt_ch, MB, 1.96 TeV, CDF-2





pt_ch, MB, 1.96 TeV, CDF-2











average pt vs nch, MB, 0.9 TeV, ATLAS





average pt vs nch, MB, 1.96 TeV, CDF-2



pt_ch, MB, 7.0 TeV, CMS



pt_ch, MB, 0.9 TeV, ATLAS



Summary

- A detailed understanding of MB interactions is especially important in very high luminosity environment (like LHC)



Summary

- A detailed understanding of MB interactions is especially important in very high luminosity environment (like LHC)
- Most important parameters of the PYTHIA MPI model are tuned



Summary

- A detailed understanding of MB interactions is especially important in very high luminosity environment (like LHC)
- Most important parameters of the PYTHIA MPI model are tuned
- Tuning results for LHC and CDF II data are inconsistent



Summary

- A detailed understanding of MB interactions is especially important in very high luminosity environment (like LHC)
- Most important parameters of the PYTHIA MPI model are tuned and compared to the AMBT1
- Tuning results for LHC and CDF II data are inconsistent
- Minimum Bias collisions are mixture of hard and soft processes, therefore, very difficult to simulate



Thanks

